



PESTICIDE PREDICAMENT: EXPLORING ENVIRONMENTAL AND HEALTH IMPACTS, AND POSSIBLE ECO-FRIENDLY SOLUTIONS

Mhaveer Singh*

Pharmacy Academy, IFTM University, Delhi Road, Lodhipur, Moradabad-244102, UP

Received on: 15.03.2024

Revised on: 22.03.2024

Accepted on: 29.03.2024

Abstract

Pesticides play a pivotal role in modern agriculture, yet their widespread use has raised significant concerns about their adverse effects on the environment and human health. This study delves into the current status of pesticide impacts, examining their repercussions on ecosystems, water sources, and the broader environment. Additionally, the potential health risks associated with pesticide exposure in humans are explored. In response to these challenges, the abstract highlights the emergence of eco-friendly management strategies, with a particular focus on bioremediation. Bioremediation, as a sustainable and nature-based approach, harnesses the power of microorganisms and plants to detoxify and degrade pesticides. The abstract emphasizes the promise of bioremediation as an effective means of mitigating pesticide pollution and fostering environmental restoration. This study encompasses the multifaceted dimensions of pesticide use, recognizing the urgency of adopting environmentally responsible practices. As society grapples with the repercussions of conventional pesticide use, the exploration of eco-friendly alternatives, especially through bioremediation, offers a ray of hope for a sustainable and harmonious coexistence with nature.

Keywords

Pesticides, Environmental Impact, Human Health, Eco-Friendly Management, Bioremediation, Sustainable Agriculture

INTRODUCTION

The Pesticide Predicament is a complex issue that encompasses environmental and health concerns, as well as the search for eco-friendly alternatives. Pesticides are chemical substances designed to control or eliminate pests that can damage crops, spread diseases, or harm livestock. While pesticides have played a crucial role in enhancing agricultural productivity, they also pose significant challenges and risks (Dayioğlu and Türker, 2021; Prabhukarthikeyan *et al.*, 2022; Soltani-Sarvestani *et al.*, 2020).

In the ever-evolving landscape of agriculture and pest management, the use of pesticides has become a cornerstone in ensuring crop yield and protecting global food supplies. However, the widespread application of these chemical agents has led to a complex and multifaceted predicament, giving rise to environmental and health concerns that demand our immediate attention. This exploration delves into the

intricate web of impacts that pesticides have on both our ecosystems and human well-being. From the contamination of soil and water to the potential health risks associated with exposure, the pesticide predicament poses a critical challenge that requires comprehensive understanding and proactive solutions. In this discourse, we navigate the intricate terrain of pesticide usage, investigating its far-reaching consequences while also delving into promising eco-friendly alternatives that hold the promise of a more sustainable and harmonious coexistence with our environment. Join us on this journey as we unravel the layers of the pesticide predicament and seek pathways toward a healthier, more environmentally conscious future (Anderson, 2017; Evans, 2018).

In the ongoing quest for sustainable environmental practices, bioremediation emerges as a powerful and innovative solution to address the challenges of pollution and contamination. The term itself suggests a biological approach

to remedying the impact of hazardous substances on our ecosystems. Bioremediation harnesses the inherent capabilities of microorganisms, plants, or enzymes to transform, degrade, or neutralize pollutants, offering a promising avenue for restoring balance to ecosystems affected by various forms of environmental degradation. This introduction aims to shed light on the principles, applications, and potential benefits of bioremediation, highlighting its role in mitigating the adverse effects of human activities on our planet and paving the way for a more sustainable and resilient future (Gaurav, 2022; Gaurav *et al.*, 2023, 2022; Gautam, 2022; Salar *et al.*, 2023)

REVIEW FINDINGS

Disastrous impact of Pesticides

The widespread use of pesticides in agriculture, while instrumental in protecting crops from pests, has also brought forth a myriad of harmful impacts on both the environment and human health. This section will explore the detrimental effects associated with pesticide use, emphasizing the need for a nuanced understanding of the trade-offs involved. From soil contamination and water pollution to the unintended harm inflicted on non-target organisms, pesticides often leave a lasting imprint on ecosystems. Furthermore, the potential health risks to humans, stemming from pesticide residues on food and exposure during application, raise significant concerns. As we unravel the harmful impacts of pesticides, it becomes evident that a balanced and sustainable approach to pest management is imperative to safeguard not only our crops but also the delicate equilibrium of the broader environment. Join us in examining the intricate web of consequences spawned by pesticide usage, prompting a critical reassessment of our agricultural practices for the well-being of both ecosystems and humanity (Tang *et al.*, 2021).

Environmental Impacts

Biodiversity Loss

The escalating use of pesticides in modern agriculture has become a major contributor to the concerning decline in global biodiversity. While these chemical agents are designed

to target specific pests, their impact often extends far beyond the intended targets, resulting in collateral damage to non-target organisms and ecosystems. This section sheds light on the profound and often overlooked consequences of pesticides on biodiversity (Saha and Bauddh, 2020).

Pesticides, designed to eliminate pests that threaten crops, can inadvertently harm beneficial insects, birds, and other wildlife in the vicinity. This unintended ecological fallout disrupts the delicate balance within ecosystems, leading to a cascade effect on various species. For example, the decline of pollinators such as bees, crucial for the reproduction of numerous plant species, can have far-reaching consequences for both flora and fauna. Moreover, pesticides can persist in the environment, accumulating in soil and water. This persistence poses a long-term threat to biodiversity as it can affect not only the immediate ecosystem but also impact interconnected habitats. Aquatic organisms, including fish and amphibians, are particularly vulnerable to waterborne pesticides, leading to disruptions in aquatic ecosystems. The loss of biodiversity due to pesticide use has ripple effects throughout the food chain, affecting predators and scavengers as well. As species disappear or decline in numbers, the resilience and adaptability of ecosystems are compromised, making them more susceptible to further environmental stressors –(Erisman *et al.*, 2016; Ollerton *et al.*, 2014; Saha and Bharadwaj, 2023; Van Der Meer *et al.*, 2020)

In recognizing the direct link between pesticide use and biodiversity loss, it becomes imperative to reassess our agricultural practices. Exploring and promoting alternative, more sustainable pest management methods is crucial to mitigate the ongoing impact on biodiversity and preserve the intricate web of life on our planet. This exploration seeks to deepen our understanding of how pesticides contribute to biodiversity decline, urging us to reconsider our approach to agriculture for the sake of a more resilient and balanced natural world. Sparrow bird and honey bees are the good examples to understand the loss of biodiversity with the use of pesticides (Figure 1).




Figure 1: Showing the yearly declined scenario of sparrow bird (Figure credit:

<https://www.theguardian.com/environment/2012/may/26/eu-farming-policies-bird-population>)

A study published on September 2019, in the journal *Science*, is the first experiment to track the effects of a neonicotinoid pesticide on birds in the wild. The study found that white-crowned sparrows who consumed small doses of an insecticide called imidacloprid suffered weight loss and delays to their migration effects that could severely harm the birds' ability to survive and reproduce. "Our study shows that this is bigger than the bees birds can also be harmed by modern neonicotinoid pesticides, which should worry us all,".

SOIL AND WATER CONTAMINATION

The pervasive use of pesticides in modern agriculture has led to a significant and widespread issue of soil and water contamination. While these chemical agents play a crucial role in protecting crops from pests, their unintended consequences on the environment, particularly on soil and water quality, pose substantial threats. This section examines the intricate ways in which pesticides contribute to contamination in soil and water systems (John and Shaik, 2015; Pretty *et al.*, 2000; Sur and Sathiavelu, 2022)

Soil Contamination

Pesticides applied to crops can accumulate in the soil, leading to long-term contamination. The persistent nature of certain pesticides means that they can remain in the soil for extended periods, negatively impacting soil health and fertility. The chemicals may leach into the deeper layers of the soil or bind to organic matter, affecting the composition of the soil microbiota (Figure 2).

This contamination can disrupt the balance of soil ecosystems, harming beneficial microorganisms and soil-dwelling organisms crucial for nutrient cycling and overall soil health. The cumulative effect of prolonged pesticide use

may result in degraded soil structure, decreased water retention capacity, and increased vulnerability to erosion.

Water Contamination

Pesticides have a propensity to move beyond the targeted crop areas through runoff, reaching nearby water bodies such as rivers, lakes, and groundwater reservoirs. This runoff introduces pesticide residues into aquatic environments, causing water contamination (Figure 2). Waterborne pesticides pose significant risks to aquatic ecosystems and the organisms that inhabit these environments.

Aquatic organisms, including fish and amphibians, can be directly affected by the presence of pesticides. Additionally, the contamination of water sources can have indirect consequences on human health when these contaminated waters are used for drinking or irrigation. The persistent and bio accumulative nature of some pesticides can lead to a gradual buildup in aquatic systems, creating a long-lasting impact on the overall water quality and the health of aquatic ecosystems. As we delve into the consequences of pesticide use on soil and water, it becomes evident that sustainable agricultural practices are crucial for mitigating these environmental impacts. Exploring alternative methods of pest control, promoting precision agriculture, and adopting integrated pest management approaches are essential steps in safeguarding the health of our soil and water systems. This exploration aims to highlight the urgent need for a balanced and environmentally conscious approach to pest management to protect these vital resources for future generations. Pesticides can leach into the soil and contaminate groundwater, posing risks to both terrestrial and aquatic ecosystems. Runoff from agricultural fields can carry pesticides into rivers and lakes, further impacting aquatic life.




Figure 2: How pesticides polluting soil and groundwater (Langenbach, 2013).

DEVELOPMENT OF RESISTANT PESTS

The development of resistant pests against pesticides is a well-documented phenomenon and a significant challenge in agriculture. Pesticides are chemicals designed to control or eliminate pests that can damage crops and reduce yields. However, over time, some pests can evolve mechanisms to survive exposure to these chemicals, leading to the emergence of pesticide-resistant populations (Figure 3). Here are some key factors contributing to the development of resistant pests (Africa, 1994; Campos *et al.*, 2016; Lengai *et al.*, 2020; Pola *et al.*, 2022).

Overuse of Pesticides

Continuous and widespread use of the same pesticides over an extended period can create strong selection pressures on pest populations. This pressure favors the survival and reproduction of individuals with natural resistance to the pesticide.

Monoculture

Large-scale monoculture, where a single crop is grown over extensive areas, provides a uniform environment for pests. This makes it easier for resistant individuals to survive and reproduce, accelerating the development of resistant populations.

Inadequate Pest Management Practices

Improper use of pesticides, such as applying sublethal doses or not following recommended application practices, can contribute to the development of resistance. Incomplete eradication of pest populations allows resistant individuals to survive and pass on their resistance to the next generation.

Genetic Variation in Pest Populations

Natural genetic variation within pest populations means that

some individuals may already have traits that confer resistance to certain pesticides. When exposed to these chemicals, only those individuals with pre-existing resistance will survive and reproduce.

High Reproductive Rates

Pests often have high reproductive rates, allowing for rapid generation turnover. This facilitates the selection and spread of resistance traits within a population in a relatively short period.

Cross-Resistance:

Some pests may develop cross-resistance, where resistance to one pesticide confers resistance to other, similar pesticides. This can happen when the same resistance mechanisms provide protection against multiple chemical compounds.

To address the issue of pesticide resistance, integrated pest management (IPM) practices are recommended. IPM involves a combination of biological, cultural, physical, and chemical control methods to minimize the reliance on pesticides and reduce the selection pressure for resistance. Crop rotation, planting pest-resistant varieties, and using alternative pest control methods are among the strategies employed in IPM to mitigate the development of resistant pests. Additionally, rotating different classes of pesticides with distinct modes of action can help slow down the development of resistance. Regular monitoring of pest populations and adapting control strategies accordingly are crucial components of effective pest management. Over time, pests can develop resistance to pesticides, leading to the need for stronger and more toxic chemicals, exacerbating the environmental impact (Siddiqui *et al.*, 2023).




Figure 3: Development of Resistant Pests




Figure 4: Health impacts associated with the exposure of synthetic pesticides (Health Risks Associated with Pesticides: <https://neem.world/health-risks-associated-pesticides/>)

Eco-Friendly Solutions of pesticide predicament

Health Impacts:

Agricultural workers and nearby communities may be exposed to pesticides through direct contact, inhalation, or ingestion. Long-term exposure has been linked to various health issues, including respiratory problems, neurological disorders, and certain cancers. Pesticide residues can persist on crops and find their way into the food supply, potentially causing harm to consumers. Chronic exposure to low levels of pesticides through the diet raises concerns about cumulative health effects (Figure 4).

Short-term exposure to high levels of pesticides can lead to acute poisoning. Symptoms may include nausea, dizziness, headaches, respiratory problems, and, in severe cases, convulsions or death. Long-term exposure to low levels of pesticides has been linked to chronic health problems. Some pesticides have been associated with the development of chronic conditions such as cancer, neurological disorders, reproductive issues, and endocrine disruption.

Some pesticides are classified as carcinogens, meaning they have the potential to cause cancer. Prolonged exposure to certain pesticides has been linked to various types of cancer, including leukemia, lymphoma, and cancers of the breast, prostate, and lung.

Certain pesticides can affect the nervous system, leading to symptoms such as dizziness, headaches, tremors, and memory loss. Prolonged exposure may contribute to the development of neurodegenerative disorders. Pesticide exposure has been associated with reproductive issues, including fertility problems, birth defects, and developmental delays in children. Pregnant women and

young children are often considered more vulnerable to the effects of pesticide exposure. Some pesticides can interfere with the endocrine system, disrupting the normal functioning of hormones. This can lead to a range of health effects, including reproductive abnormalities, metabolic disorders, and developmental issues.

Inhalation of pesticide residues or vapors can cause respiratory problems, including irritation of the nose and throat, coughing, and difficulty breathing. Certain individuals may be allergic to specific pesticides, leading to skin rashes, itching, or respiratory symptoms upon exposure. Farmworkers and individuals involved in the production, application, or handling of pesticides face a higher risk of exposure and associated health effects. Proper protective measures and adherence to safety guidelines are crucial to minimizing these risks. Pesticides can leach into groundwater, posing a risk to drinking water supplies. Prolonged exposure to contaminated water sources can lead to chronic health issues.

To mitigate the harmful health impacts of pesticides, it is important to implement proper safety measures, use protective equipment, follow recommended application practices, and explore alternative pest control methods. Additionally, regulatory agencies play a crucial role in setting and enforcing safety standards for the use of pesticides in agriculture and other industries. Public awareness and education about the potential risks of pesticide exposure are also essential for promoting safer practices and minimizing health risks.

INTEGRATED PEST MANAGEMENT (IPM):

Integrated Pest Management (IPM) is a holistic and sustainable approach to managing pests that aims to minimize economic, environmental, and health risks. IPM combines various strategies to control pests effectively while reducing the reliance on chemical pesticides. The goal is to maintain pest populations at levels below those causing economic or aesthetic damage while promoting natural pest control mechanisms. Here are the key components of Integrated Pest Management (Barzman *et al.*, 2015; Chandler *et al.*, 2011; Fahad *et al.*, 2021; Khan *et al.*, 2021; Pretty and Bharucha, 2015).

Biological Control and Cultural Controls

Encouraging natural predators, parasites, and pathogens that feed on or parasitize pests. This can include releasing beneficial organisms like ladybugs, predatory mites, or using pathogens like *Bacillus thuringiensis* (Bt) for targeted pest control. Modifying agricultural or landscaping practices to make the environment less favorable for pests. This can involve crop rotation, planting resistant varieties, adjusting planting dates, and promoting diverse cropping systems.

Mechanical and Physical Controls

Using physical barriers, traps, or other mechanical methods to prevent pests from reaching crops. Examples include using row covers, sticky traps, and employing cultivation practices to disrupt pest life cycles.

Chemical Controls

Judicious use of pesticides when necessary. In IPM, chemical controls are considered a last resort and are applied selectively and in accordance with a well-defined strategy. It involves choosing pesticides with lower environmental impact, rotating chemical classes, and using them at the right time and in the right amount.

Monitoring, Scouting and Cultural Practices

Regularly assessing pest populations and their impact on crops. This involves monitoring pest levels, identifying beneficial organisms, and using thresholds to determine when control measures are necessary. Implementing practices that promote a healthy and robust crop, making it less susceptible to pest damage. This can include proper irrigation, fertilization, and maintaining optimal plant spacing.

Host Plant Resistance

Developing and using crop varieties that are resistant or tolerant to pests. This can reduce the need for chemical controls and support a more sustainable approach to pest management.

Public Awareness and Education

Educating farmers, landscapers, and the public about the principles of IPM, the importance of biodiversity, and the potential risks associated with excessive pesticide use. This helps foster a broader understanding of sustainable pest management practices.




Figure 5: Different steps involved in Integrated Pest Management process.

Economic Thresholds

Establishing economic thresholds to determine when the cost of controlling pests is justified by the potential damage they can cause. This helps in making informed decisions about when and how to implement control measures.

Adaptive Management

Continuously assessing and adjusting pest management strategies based on monitoring and evaluation. This involves learning from experience and adapting practices to changes in pest populations or environmental conditions.

By integrating these various components, IPM provides a comprehensive and environmentally friendly approach to pest management that seeks to balance the need for crop protection with long-term sustainability. It has gained widespread acceptance as an effective and responsible approach in agriculture, horticulture, and landscaping. IPM focuses on a holistic approach to pest control, incorporating biological, cultural, and mechanical methods along with judicious use of pesticides. This approach minimizes the environmental impact and reduces the reliance on chemical treatments.

ORGANIC FARMING PRACTICES

Organic farming is an agricultural approach that emphasizes sustainability, soil health, and the use of natural methods to grow crops and raise livestock. Organic farming practices aim to minimize the use of synthetic inputs, such as chemical fertilizers and pesticides, and promote ecological balance. Here are some key principles and practices associated with organic farming (2023; Atoma *et al.*, 2020; Galnaitytė *et al.*, 2017; Kanagasabapathi and Sakthivel, 2019; Kavaskar and Govind, 2020; Kumar *et al.*, 2022).

Crop Rotation

Organic farmers often use crop rotation to promote soil fertility and reduce the risk of pests and diseases. Rotating different crops on the same land over time helps break pest and disease cycles and enhances soil structure.

Composting

Composting is a common practice in organic farming for

recycling organic matter. Compost, which is a mixture of decomposed plant and animal materials, is used to improve soil structure, fertility, and water retention.

Cover Cropping

Growing cover crops, such as legumes, on fallow fields or between cash crops helps prevent soil erosion, suppress weeds, and improve soil fertility by adding organic matter. Cover crops also provide habitat for beneficial insects.

Organic Fertilizers: Organic farmers use natural sources of nutrients, such as compost, manure, and green manure, to fertilize their crops. These materials release nutrients slowly, promoting long-term soil health.

Biological Pest Control: Instead of relying on synthetic pesticides, organic farmers employ biological control methods. This includes introducing or attracting natural predators, using beneficial insects, and implementing other measures to manage pest populations.

Crop Diversity: Growing a variety of crops helps break pest cycles, reduces the risk of diseases, and promotes overall ecosystem health. Crop diversity can also enhance soil fertility and nutrient cycling.

No Synthetic Chemicals: Organic farming prohibits the use of synthetic chemicals, including synthetic pesticides and fertilizers. Instead, farmers rely on natural alternatives and sustainable practices to manage pests and enhance soil fertility.

Livestock Integration: Organic farms often integrate livestock into the farming system. Livestock contribute to nutrient cycling through manure, and their presence can be beneficial for pest control and weed management.

Non-GMO (Genetically Modified Organisms): Organic farming standards typically prohibit the use of genetically modified organisms (GMOs). Organic farmers use traditional breeding methods to develop crop varieties with desirable traits.

Sustainable Water Management: Organic farmers prioritize efficient water use and often employ water conservation practices, such as mulching and drip irrigation, to reduce water consumption.

Minimal Soil Disturbance: Organic farming emphasizes minimal soil disturbance to preserve soil structure and minimize erosion. Practices like no-till or reduced-till farming are common in organic systems.

Certification and Standards: To be recognized as organic, farms must adhere to specific standards and undergo certification processes established by organic certifying bodies. These standards ensure that organic practices are followed, and consumers can trust the organic label.

Organic farming practices aim to create a sustainable and

environmentally friendly system that prioritizes soil health, biodiversity, and the well-being of both the environment and consumers. While organic farming has gained popularity, it's important to note that challenges exist, including potential lower yields in certain cases and the need for careful management to control pests and diseases without synthetic inputs. Organic farming avoids synthetic pesticides and promotes the use of natural alternatives, such as neem oil, diatomaceous earth, and beneficial insects. This approach aims to minimize environmental impact and produce healthier, pesticide-free food.




Figure 6: Different aspects of organic farming (<https://www.zauca.com/organic-farming-website-design-rs-5900-organic-farm-website-development-company-near-me/>).

AGROECOLOGY

Agroecology is an interdisciplinary approach to agriculture that integrates ecological principles into farming systems. It emphasizes the development of sustainable and resilient food production systems while minimizing the use of external inputs like synthetic pesticides and fertilizers. Agroecology seeks to promote ecological balance, enhance biodiversity, and improve the overall health of agricultural ecosystems. Here are key principles and components associated with agroecology (Duff *et al.*, 2022; Kerr *et al.*, 2022; López-García and de Molina, 2021; van der Ploeg, 2021).

Biodiversity

Agroecology encourages the cultivation of diverse crops and the promotion of on-farm biodiversity. This diversity helps enhance ecosystem resilience, reduce the risk of pests and diseases, and improve overall ecosystem health.

Crop Rotation and Polyculture

Rotating crops and planting multiple crops together (polyculture) are common practices in agroecology. These techniques disrupt pest and disease cycles, improve soil

fertility, and contribute to a more balanced ecosystem. Integrating trees and shrubs into agricultural landscapes is a key aspect of agroecology. Agroforestry systems provide multiple benefits, such as improved soil structure, increased biodiversity, carbon sequestration, and additional income sources through the production of fruits, nuts, or timber. Minimizing soil disturbance through practices like no-till or reduced tillage helps maintain soil structure, reduce erosion, and enhance water retention. This is crucial for sustaining soil health and fertility.

Efficient water management practices, such as rainwater harvesting, drip irrigation, and mulching, are emphasized in agroecology. These practices help conserve water resources and reduce the environmental impact of agriculture. Agroecology values traditional knowledge and local farming practices. Indigenous and traditional farming systems often hold valuable insights into sustainable agricultural practices that can be incorporated into modern agroecological approaches. Agroecology emphasizes the use of natural enemies, biological control methods, and other ecological processes to manage pests. This reduces the reliance on synthetic pesticides and promotes a healthier balance between pests and their natural predators —(Abbona et al., 2007; Bocco et al., 2019; Costa-Pierce et al., 2011; Tully and Ryals, 2017).

Maintaining and improving soil health is a core principle of agroecology. Practices such as cover cropping, composting, and organic matter incorporation contribute to soil fertility, structure, and nutrient cycling. Agroecology takes into account the social and economic dimensions of agriculture. It aims to create equitable and sustainable food systems that benefit local communities, support small-scale farmers, and enhance food security.

Agroecology often involves a participatory and community-based approach, where farmers actively engage in decision-making processes. Local knowledge and community involvement are considered essential for the success of agroecological initiatives. Agroecological practices are designed to build resilience in farming systems against the impacts of climate change. Diverse and resilient ecosystems are better equipped to adapt to changing environmental conditions. Agroecology promotes education and training at various levels, encouraging farmers to understand the ecological principles underlying their practices and fostering a culture of continuous learning.

Agroecology represents a shift toward more sustainable, regenerative, and ecologically conscious agricultural practices. It aligns with the broader goals of sustainable development, emphasizing the interconnectedness of ecological, social, and economic factors in agriculture. Many believe that agroecology can contribute to addressing global challenges such as food insecurity, environmental degradation, and climate change. Agroecological practices focus on sustainable farming systems that enhance biodiversity, soil health, and resilience to pests and diseases (Abbona et al., 2007; Tully and Ryals, 2017).

Furthermore, continued research into alternative pest control methods, such as biopesticides and precision agriculture, can contribute to more sustainable and effective solutions. However, addressing the pesticide predicament requires a comprehensive and multi-faceted approach that balances the need for pest control with environmental and health considerations. Transitioning towards eco-friendly alternatives and promoting sustainable agricultural practices are essential steps in mitigating the negative impacts of pesticides on our ecosystems and well-being.

Conclusion

In conclusion, the pesticide predicament poses a multifaceted challenge, intertwining environmental and health impacts that demand urgent attention and innovative solutions. The indiscriminate use of chemical pesticides has undeniably led to severe ecological imbalances, adversely affecting biodiversity, soil health, and water quality. Simultaneously, the detrimental consequences extend to human health, with potential long-term risks linked to pesticide exposure. However, amidst this predicament lies the opportunity for transformative change. The exploration of eco-friendly alternatives emerges as a promising avenue to mitigate the adverse effects of conventional pesticides. Embracing sustainable agricultural practices, integrated pest management, and the development of biopesticides can pave the way towards a harmonious coexistence of agriculture and the environment. Public awareness and education are essential components in fostering a collective commitment to environmentally responsible practices. Governments, researchers, and stakeholders must collaborate to enact and enforce stringent regulations that prioritize the development and adoption of eco-friendly solutions. As we navigate the pesticide predicament, it is imperative to recognize the interconnectedness of environmental and human well-being, striving for a future where agricultural practices coexist harmoniously with nature, ensuring a sustainable and healthier planet for generations to come.

Acknowledgement

The author would like to thank Pharmacy Academy, IFTM University, Delhi Road, Lodhipur, Moradabad-244102, UP for providing continuous support to complete the study.

Conflict of interest

Author declares no conflict of interest

References

1. **Prabhukarthikeyan SR, Keerthana U, Baite MS, Panneerselvam P, Mitra D, Naveen Kumar, Parameswaran C, Cayalvizhi B, Muthu Kumar A, Harish S, Rath PC.** (2022). Bacillus rhizobacteria: A versatile biostimulant for sustainable agriculture. *New and Future Developments in Microbial Biotechnology and Bioengineering*, 33-44.
2. **Dayioğlu MA, Türker U.** (2021). Digital transformation for sustainable future-agriculture 4.0: A review. *Journal of Agricultural Sciences (Tarım Bilimleri Dergisi)*, 27(4), 373-399.

3. **Soltani-Sarvestani M, Shamspur T, Mostafavi A, Falahati-Pour SK, Mohamadi M, Saljooqi A.** (2020). A novel preconcentration/extraction method for fenitrothion residues and its electrochemical sensing in pistachio kernels. *Journal of Materials Science: Materials in Electronics*, 31, 11268-11275.
4. **Anderson JM.** (2017). The Annihilation of nature: Human extinction of birds and mammals. *Transactions of the Royal Society of South Africa*, 72(3), 303–304.
5. **Evans L.** (2018). Bioremediation techniques for agricultural pollutants. Reference Module in Earth Systems and Environmental Sciences, 1-5.
6. **Sharma GI, Khan MU, Zahiruddin S, Basist P, Ahmad S.** (2023). Multi-mechanistic and therapeutic exploration of nephroprotective effect of traditional ayurvedic polyherbal formulation using in silico, in vitro and in vivo approaches. *Biomedicines*, 11(1), 168-188.
7. **Gaurav, Khan MU, Basist P, Zahiruddin S, Ibrahim M, Parveen R, Ibrahim M, Parveen R, Krishnan A, Ahmad S.** (2022). Nephroprotective potential of Boerhaavia diffusa and Tinospora cordifolia herbal combination against diclofenac induced nephrotoxicity. *South African Journal of Botany*, 151(B), 238-247.
8. **Salar S, Gaurav, Sharma P.** (2023). Quality Control and multi-targeted therapeutic approach of nycanthes arbor-tristis for management of hepatic disease and associated complications. *Pharmacognosy Magazine*, doi:10.1177/09731296231189619.
9. **Gaurav.** (2022). GC–MS metabolomics and network pharmacology-based investigation of molecular mechanism of identified metabolites from Tinospora cordifolia (Willd.) miers for the treatment of kidney diseases. *Pharmacognosy Magazine*, 18, 548-558.
10. **Gautam G.** (2022). Network pharmacology-based validation of traditional therapeutic claim of momordica charantiain alleviating diabetic nephropathy. *Journal of CAM Research Progress*, 1, 1-10.
11. **Saha L, Baudhh K.** (2020). Sustainable agricultural approaches for enhanced crop productivity, better soil health, and improved ecosystem services. *Ecological and Practical Applications for Sustainable Agriculture*, 1-23.
12. **Ollerton J, Erenler H, Edwards M, Crockett R.** (2014). Extinctions of aculeate pollinators in Britain and the role of large-scale agricultural changes. *Science*, 346(6215), 1360-1362.
13. **Erisman JW, van Eekeren N, de Wit J, Koopmans C, Cuijpers W, Oerlemans N, Koks BJ.** (2016). Agriculture and biodiversity: A better balance benefits both. *AIMS Agriculture and Food*, 1(2), 157-174.
14. **Saha S, Bharadwaj A.** (2023). A step towards smart agriculture using metallic nanostructures. *Plant Stress*, 10, 100216.
15. **Van Der Meer M, Kay S, Lüscher G, Jeanneret P.** (2020). What evidence exists on the impact of agricultural practices in fruit orchards on biodiversity? A systematic map. *Environmental Evidence*, 9, 2. <https://doi.org/10.1186/s13750-020-0185-z>
16. **Sur S, Sathivelu M.** (2022). A concise overview on pesticide detection and degradation strategies. *Environmental Pollutants and Bioavailability*, 34(1), 112-126.
17. **John EM, Shaik JM.** (2015). Chlorpyrifos: pollution and remediation. *Environmental Chemistry Letters*, 13, 269-291.
18. **Pretty JN, Brett C, Gee D, Hine RE, Mason CF, Morison JIL, Raven H, Rayment MD, Van der Vijn G.** (2000). An assessment of the total external costs of UK agriculture. *Agricultural Systems*, 65(2), 113-136.
19. **Langenbach T.** (2013). Persistence and bioaccumulation of persistent organic pollutants (POPs). *Applied Bioremediation - Active and Passive Approaches*, 13. <http://dx.doi.org/10.5772/56418>
20. **Lengai GMW, Muthomi JW, Mbega ER.** (2020). Phytochemical activity and role of botanical pesticides in pest management for sustainable agricultural crop production. *Scientific African*, 7, e00239. <https://doi.org/10.1016/j.sciaf..e00239>
21. **Campos EVR, de Oliveira JL, Pascoli M, de Lima R, Fraceto LF.** (2016). Neem oil and crop protection: From now to the future. *Frontiers in Plant Science*, 7, 1494. <https://doi.org/10.3389/fpls.2016.01494>
22. **Pola S, Kesharwani AK, Singh J, Singh D, Kalia VK.** (2022). Endophytic ability of indigenous Bacillus thuringiensis strain VKK-BB2: new horizons for the development of novel insect pest-resistant crops. *Egyptian Journal of Biological Pest Control*, 32, 8-20. <https://doi.org/10.1186/s41938-022-00512-y>
23. **Africa W.** (1994). Development of Multiple Pest Resistant Crop Cultivars. *Journal of Agricultural Entomology*, 11(3), 225-253.
24. **Barzman M, Bårberi P, Birch ANE, Boonekamp P, Dachbrodt-Saaydeh S, Graf B, Hommel B, Jensen JE, Kiss J, Kudsk P, Lamichhane JR, Messéan A, Moonen AC, Ratnadass A, Ricci P, Sarah JL, Sattin M.** (2015). Eight principles of integrated pest management. *Agronomy for Sustainable Development*, 35, 1199-1215.
25. **Fahad S, Saud S, Akhter A, Bajwa AA, Hassan S, Battaglia M, Adnan M, Wahid F, Datta R, Babur E, Danish S, Zareit, Irshad I.** (2021). Bio-based integrated pest management in rice: An agroecosystems friendly approach for agricultural sustainability. *Journal of the Saudi Society of Agricultural Sciences*, 20(2), 94-102.

26. **Pretty J, Bharucha ZP.** (2015). Integrated pest management for sustainable intensification of agriculture in Asia and Africa. *Insects*, 6(1), 152-182.
27. **Khan FZA, Manzoor SA, Gul HT, Ali M, Bashir MA, Akmal M, Haseeb M, Imran MU, Taqi M, Manzoor SA, Lukac M, Joseph SV.** (2021). Drivers of farmers' intention to adopt integrated pest management: a case study of vegetable farmers in Pakistan. *Ecosphere*, (10), e03812. <https://doi.org/10.1002/ecs2.3812>
28. **Chandler D, Bailey AS, Mark Tatchell G, Davidson G, Greaves J, Grant WP.** (2011). The development, regulation and use of biopesticides for integrated pest management. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 366(1573), 1987-1998.
29. **Kumar A, Singh K, Verma P, Singh O, Panwar A, Singh T, Kumar Y, Raliya R.** (2022). Effect of nitrogen and zinc nanofertilizer with the organic farming practices on cereal and oil seed crops. *Scientific Reports*, 12, 6938. <https://doi.org/10.1038/s41598-022-10843-3>
30. **Badu BR.** (2023). Organic Farming Practices For Horticultural Sustainability. *International Journal for Multidisciplinary Research*. 5(4). <https://doi.org/10.36948/ijfmr.2023.v05i04.5367>
31. **Atoma NC, Anyoha PN, Chikaire UJ, Uyoyuo S.** (2020). Adoption of organic farming practices in the rural household of South-South, Nigeria—a case study. *Organic Agriculture*, 10, 173–178.
32. **Kavaskar M, Govind S.** (2020). Factors influencing symbolic adoption behaviour of paddy growers on organic farming practices. *Plant Archives*, 20(1), 2956-2960.
33. **Galnaitytė A, Kriščiukaitienė I, Baležentis T, Namiotko V.** (2017). Evaluation of technological, economic and social indicators of different farming practices in Lithuania. *Economics & Sociology*. 10(4), 189-202.
34. **Kanagasabapathi K, Sakthivel V.** (2019). An analysis of adoption of organic farming practices of tribal farmers of Kolli hills. *Plant Archives*, 19(2), 3533-3534.
35. **van der Ploeg JD.** (2021). The political economy of agroecology. *The Journal of Peasant Studies*, 48(2), 274-297.
36. **Kerr RB, Liebert J, Kansanga M, Kpienbaareh D.** (2022). Human and social values in agroecology: A review. *Elementa: Science of the Anthropocene*, 10 (1), 00090.
37. **Duff H, Hegedus PB, Loewen S, Bass T, Maxwell BD.** (2022). Precision agroecology. *Sustainability*. 2022, 14(1), 106-123.
38. **López-García D, de Molina MG.** (2021). An operational approach to agroecology-based local agri-food systems. *Sustainability*, 13(15), 8443.
39. **Tully K, Ryals R.** (2017). Nutrient cycling in agroecosystems: Balancing food and environmental objectives. *Agroecology and Sustainable Food Systems*, 41(7), 761-798.
40. **Abbona EA, Sarandón SJ, Marasas ME, Astier M.** (2007). Ecological sustainability evaluation of traditional management in different vineyard systems in Berisso, Argentina. *Agriculture, Ecosystems & Environment*, 119(3-4), 335-345.
41. **Bocco G, Solis Castillo B, Orozco-Ramirez Q, Ortega-Iturriaga A.** (2019). Terraced agriculture in the adaptation to climatic variability in the Mixteca Alta, Oaxaca, Mexico. *Journal of Latin American geography*, 18(1), 141-168.
42. **Costa-Pierce B, Bartley D, Hasan M, Yusoff F, Kaushik S, Rana K, Lemos D, Bueno P, Yakupitiyage A.** (2010). Responsible use of resources for sustainable aquaculture. *Environmental Science, Agricultural and Food Sciences*. ID: 82669991 <https://api.semanticscholar.org/CorpusID:82669991>